

Figure 1

Class No.....

[illegible]

SVEN HEDIN

SCIENTIFIC RESULTS

OF A JOURNEY IN

CENTRAL ASIA


1899—1902

VOL VI. PART II

G E O L O G Y

BY

DR HELGE BÄCKSTRÖM AND HARALD JOHANSSON



STOCKHOLM

LITHOGRAPHIC INSTITUTE OF THE GENERAL STAFF
OF THE SWEDISH ARMY

STOCKHOLM

KUNGL. BOKTRYCKERIET. P. A. NORSTEDT & SÖNER

1907

GEOLOGY

BY

HELGE BÄCKSTRÖM AND HARALD JOHANSSON.

Dr. SVEN HEDIN's geological collections from this voyage consist of not less than 800 rock specimens, with few exceptions taken from bed rocks in different places all along his route. Through the kindness of Dr. HEDIN these large collections were presented to the geological department of the University of Stockholm, where they are now deposited, together with the numerous slides for study made from them, and the specimens collected by Dr. HEDIN in the same region in 1896 which previously had also been presented to the University of Stockholm.

The whole material was studied by us, the rocks determined, the descriptive catalogue of the specimens given below worked out, and finally an effort was made to construct a geological map of the area. This map, naturally, could for the most part become nothing more than a geological map of the route, still we hope that it may serve as a skeleton-map to be filled out by future explorations.

The rocks collected are classified on the map as follows:

- Red sandstone series,
- Greywacke series,
- Slate and Phyllite series,
- Gneiss and Mica Schist series,
- Dense Limestone,
- Crystalline Limestone,
- Greenstones and Greenstone-schists.
- Young volcanic rocks,
- Old volcanic rocks,
- Granites and related deep-seated igneous rocks.

Red Sandstone series.

The deposits belonging to this series consist of sandstone and sandy slates containing here and there layers of gypsum and also conglomerates. The rocks generally exhibit red or violet colours, although greenish or gray rocks have also been included. According to HEDIN the soils in the sandstone regions are bright red in colour.

The sandstones consist of quartz grains which are generally angular and which are embedded in a matrix very red in colour and generally free from calcite. In this series, however, there have also been included some calcareous, loose, non-metamorphosed sandstones or fine sandy slates usually with grayish colours. (Ex. A: 315—319; B: 113—114.)

The conglomerates are generally very polymict, with well rounded pebbles of dense limestones, dark slates, greenish greywackes or quartz; in several places also of porphyries. (Ex. A: 122; B: 100.)

The sandstones are not dynamo-metamorphic; they have taken part in the formation of the mountain ranges but the dips are often flat. — The series has a considerable extension in Eastern Tibet, probably even greater than the map shows, since the specimens on which the map is based were collected along the route, i. e. in the valleys and passes, and these specimens sometimes consist of other rocks, although, according to HEDIN's description, the surrounding mountains are made up of sandstone. As mentioned above there are also large districts in which the bed rock does not appear although from the reddish colour of the soil it is evident that it is sandstone.

Greywacke series.

This name has been used for a widely distributed and rather constant type of sedimentary rocks. The rocks in question generally show dark greenish-grey or brownish-green colours and are sometimes finely rust-spotted. They are composed of fragments of quartz, felspar and light mica in a matrix, generally rich in calcite; very often also fragments of black slates and dense limestones are found. The more coarse-grained varieties consist of a lightly weathered granitic material with some detritus of slate (Ex. A: 309, 321, 323, 27, 28). Real conglomerates apparently do not occur. — The most common types are constituted by more fine-grained angular detritus with larger grains of felspar, quartz or mica sprinkled in without marked arrangement or layering. In a few cases the quartz grains predominate, thus producing varieties which have been referred to in the catalogue as greywacke-sandstones and quartzite-sandstones. Besides the greywackes mentioned above in which the grains are distinguishable macroscopically, more fine-grained microclastic rocks, "greywacke-slates" have a wide distribution. They occur, according to the descriptions given by HEDIN in his work on his journey of 1896, in alternate bedding with the coarser varieties. Their color is generally a dark grey, caused by the presence of a carbonaceous pigment. In these greywacke-slates

there may sometimes be seen twig-like accumulations of apparently carbonaceous matter which may possibly be interpreted as being fossil plants. (Ex. A: 312, 112, 170 a. o.) These greywacke-slates are harder than true clay slates and do not show well pronounced slaty cleavage but more uneven fracture. — The clastic structure is microscopically well preserved and the rocks do not show any considerable recrystallisation.

The greywackes are everywhere strongly dislocated, showing steep dips. They show very marked signs of pressure, are very much hardened, often show slickensides as well as fissures filled up with carbonates or quartz. In certain regions they are rich in quartz nodules produced by secretion (Ex. B: 274—276, a, h etc.); most of the numerous quartz and quartzite specimens in the collection probably came from such secretions. In some parts the schistosity is specially developed, thus producing glistening separation plans; the greenish greywackes here pass over into greenish schists with parallel jointing (A: 223, 88 a. o.); the dark greywackes into darkly glistening hard schists, often rust-spotted (Ex. B: 275); the dark more fine-grained greywacke-slates pass over into phyllitic schists which it has probably not been possible to separate from certain phyllites, not belonging to the greywacke series.

Some very much hardened quartzite-sandstones of rather different type have also been mapped as greywacke.

The greywacke series is markedly older than the red sandstones but younger than, at least, part of the schists and limestones; probably also younger than most of the granites.

Representatives for this series have been found over the whole region covered by the route, from the beginning to the end.

Clay slates and phyllites.

These rocks are found here and there; are, however, wanting in the north-east region, which is comparatively most crystalline. Some of them appear to be geologically related to the dark, dense limestones, others are probably related to the above mentioned greywacke-slates. — The same color has also been used on the map for some specimens of black siliceous schists (Ex. A: 192, 196; B: 41, 74, 75). — In the neighbourhood of Ladak a series of rather varying schists have been mapped as phyllites.

On the border of the great granite regions several examples of contact metamorphism are found, as spotted slates (A: 30) and hornfelses.

Dense limestones and dolomites.

Of these varying types occur, all of which can probably not be referred to a single geological formation, although most of them appear to be older than not only the red sandstones but also than the greywackes and a good many of the

granites. It is possible to distinguish to a certain extent a series of lighter, purer limestones (with grey, pink, yellowish etc. colors), occasionally with chert concretions (Ex. B: 81, 116, 121), and a series of dark, more impure bituminous limestones, passing over into calcareous clay slates. — Judging from the dips of the strata, the limestones are everywhere dislocated, generally also crushed and filled with calcite veins. Semicrystalline or microcrystalline types are often found. Near the great mass of granite in NE Tibet dark limestones and calcareous clay slates are found, containing small crystals of scapolite and diopside, also lumps of magnetic pyrites (Ex. A: 273, 271), a specimen of more crystalline calcite-diopside-rock (A: 272) also comes from here. These rocks apparently are due to the metamorphic action of the granite.

The limestones are most richly represented in the south eastern parts of the route and there they are also least metamorphosed. From this region there have also been brought several specimens containing remains of fossils. Unfortunately all these traces of fossils are indeterminable, on account of which the most interesting question of the geological age of these Tibetan deposits cannot be solved by this material. But as it appears to be indubitable that the question of the age of the deposits might easily be solved if specimens were collected here with special regard to the fossil remains which they contain, the localities from which we have noted traces of fossils in the limestone-specimens have been marked with a blue **F** on the map and an enumeration of the localities is here given, which may be of service in future investigations:

A-series:

1900. $13/9$ No. 142.

B-series:

1901.	$10/7$	No. 78
	$21/7$	89
	$24/8$	96
		98
	$25/8$	103
	$10/9$	120
	$11/9$	121

1901.	$11/9$	No. 124
	$14/9$	125
		126
	$15/9$	127
	$18/9$	128
	$25/9$	129—131
	$10/10$	158.

Crystalline limestones and dolomites.

Some of the crystalline limestones are found in close connection with the dense varieties, others in the more highly metamorphosed districts together with gneisses and mica schists. As dark bituminous veins are sometimes found and also passages into semicrystalline, dark dense limestones, it seems rather probable that the crystalline varieties, genetically considered, are only more intensely metamorphosed facies of the dense varieties. — Special attention must be drawn to the many dolomites found.

Gneiss and Mica Schist.

In this series have been included various strongly crystalline types of schists: gneisses, granulites, mica schists, hornfels-like rocks etc. The occurrence of such rocks is surprisingly small in view of the region being mountainous; they are completely absent in the southern two thirds of the region covered by the map, and are only found to any extent in the north.

Schistose, gneiss-like granites have been mapped as granites as far as possible.

The gneisses are essentially grey biotite- or biotite-hornblende-bearing types rich in white felspar. Of other types there may be mentioned:

A: 364, 368, 373 = light granulitic rocks.

A: 369, 376 = garnet-bearing gneisses.

B: 19 = pyroxene-gneiss with calcite and scapolite.

Mica schists are rather rare. — A: 370 is a coarse-grained garnet-bearing variety; A: 363 a fine-grained muscovite-chlorite-schist; etc.

In this section there have also been included a series of close-grained, dark grey or violet grey hornfelslike rocks, distinguished by a very strongly coloured red-brown biotite, and in part rather rich in felspar. Some of them seem to be partly recrystallized greywackes, others are evidently crystallized phyllites; these rocks apparently have got their characters by contact-metamorphism from the great masses of granite and their extensions.

Greenstones and Greenstone-schists.

Diabases, comparatively unaffected by pressure, are sometimes found; in some cases as dykes, in other cases their geological mode of occurrence is not noted. The diabases and connected "greenstones" are, however, in general more or less completely converted into amphibolites or very schistose "greenstone-schists". Especially is this the case in the most north-eastern part of the route, where, on the whole, the metamorphism has been most intense. From here also a few schistose gabbros were brought home.

Granites.

Granites frequently occur in the north-eastern parts as well as in the district north of Ladak; on the other hand they are lacking in the whole of the southern region. Of the three masses north of Ladak, the most northern and the most southern consist of soda-granites, while the middle one is made up of quartz-dioritic rocks. — In the great granite-region in north-eastern Tibet granites of very varying types occur, strongly modified by pressure as well as comparatively little changed. As mentioned above some of them border upon schists which are evidently contact-metamorphosed and are consequently younger than part of the schist-formations; on the other hand the felspathic material of the greywackes indicates a derivation from older granites.

Effusive igneous rocks.

It seems possible to carry out in this region the customary division of these rocks into an older and a younger series, the latter characterized by a preserved glassy groundmass as well as by a greater freshness of the constituent minerals. The former has been referred to on the map as "*old volcanic rocks*", the latter as "*young volcanic rocks*".

Effusive igneous rocks of older type are found here and there along the route; their principal distribution, however, is in NE Tibet (Nos. A: 51—64, 245—248, 250). Here occurs a rich variety of quartzporphyrytuffs, some composed of the finer detritals and showing the characteristic ash-structure, others are again made up of crystals of quartz and feldspar and of angular pieces of quartzporphyries of different types; subordinately also of detritals from dark porphyritic rocks. In the western part of the region, the porphyritic character becomes more accentuated. — The most of these tuffs and associated porphyries do not exhibit marked signs of strain, but on the northern border examples of dynamometamorphic action occur. Such schistose quartz-porphyries or quartz-porphyrates, passing into sericite-schists, are met with on several places along the route.

Young volcanic rocks were also met with during HEDIN's previous journey at those points along this his route, which are indicated on the map. The specimens brought home then were studied by one of us.* This investigation showed that the rocks were a *bronzite-andesite*, characterized by the fact that all the crystallized pyroxene is rhombic, while the monoclinic pyroxene is completely or almost entirely lacking.

To a similar type belong also the specimens collected on the new journey from the localities at 35° N. L., 89° E. Gr. (A: 144—152) and 36° 15' N. L., 88° 30' E. Gr. (B: 46); still varieties which besides the dominating bronzite contain not inconsiderable quantities of monoclinic pyroxene were also found here.

An interesting holocrystalline variety of this type of rock occurs between Naktsong-tso and Selling-tso, the most southern point on HEDIN's route (B: 123). Macroscopically examined the rock shows a close-grained violet-brown groundmass with small phenocrysts of fresh and glistening pyroxene. This pyroxene seen under the microscope is found to be exclusively rhombic bronzite; only on the borders of the larger crystals, as well as here and there among the pyroxenes of the groundmass, the bright interference colours indicate the presence of small quantities of monoclinic pyroxene. Associated with the bronzite and distinctly younger than this, brown biotite occurs, but never as phenocrysts. The groundmass consists — excepting rhombic and some monoclinic pyroxene, biotite, ores etc. — essentially of lathshaped plagioclase with almost trachyoidal appearance. — About the mode of occurrence of this rock nothing is said in HEDIN's notes.

* H. BACKSTRÖM, Über jungvulkanische Eruptivgesteine aus Tibet. Petermanns Mittheilungen, Ergänzungsheft 131, p. 375 (1900).

From $35^{\circ}25'$ N. L., 90° E. Gr. a different type was brought home (A: 107): an andesite rich in phenocrysts of plagioclase and a dark mineral, now totally decomposed to greenish blue hornblende and chlorite. This rock formed two small conical knobs (Scientific Results III p. 103).

At Camp XLV ($34^{\circ}40'$ N. L., $89^{\circ}50'$ E. Gr.) there occurs a dark biotite-bearing hornblende-andesite, quite undecomposed and with glassy groundmass; another specimen shows the same rock weathered and with lighter colour (A: 130, 131).

No. A: 121 ($34^{\circ}45'$ N. L., $90^{\circ}20'$ E. Gr.), a hard, violet-black rock (Scientific Results III p. 129) is a rather weathered holocrystalline trachytoidic andesite; A: 126 ($34^{\circ}45'$ N. L., 90° E. Gr.) a hard and compact andesitic tuff. — No. A: 135 ($34^{\circ}35'$ N. L., $89^{\circ}35'$ E. Gr.) is a compact, somewhat weathered hornblende-andesite of a different type.

At $31^{\circ}40'$ N. L., $86^{\circ}49'$ E. Gr. LITTLEDALE (*vide* footnote on next page) had reported a volcano called "Tongo", which HEDIN visited on Oct. 4th 1901 and declared not to be a volcano (Sc. Res. IV p. 125—131). He found there as bed rocks a conglomerate, belonging to the red sandstone series, and weathered diabases; he also collected some boulders, apparently derived from the upper parts of the mountain. Among these there are weathered porphyrites, light yellow or white, compact, holocrystalline-granophyric, evidently rather young rocks, and finally also "a kind of light green porphyry or granite", which showed itself to be a rather weathered hornblende-andesite.

Outside eastern Tibet young volcanic rocks have been brought home from the shore of Oman-tso ($32^{\circ}25'$ N. L., $82^{\circ}45'$ E. Gr.; Scientific Results IV p. 196; Number of specimens: B: 179—187.) — Some of the rocks from here are rhyolitic tuffs, which, according to HEDIN's diary, show very distinct bedding; the rock "shows long, red, sharply defined edges, jutting out of the detritus on the southern flank of the mountains". — Under the microscope these tuffs show the characteristic ash-structure admirably well developed. Except as tuffs rhyolites do not seem to occur here; on the other hand several specimens consist of hard compact trachytes. They are violet-grey or red, sometimes with secretions of agate, and show under the microscope phenocrysts of biotite and an anorthoclastic feldspar in a groundmass with trachyoidal structure.

Far up in the north-east, between Camps CXLI and II, there is a small district of rather weathered but apparently not very old volcanic rocks, partly basic andesites, partly trachytes. (A: 412 etc.)

About the date of the volcanic activity in these regions some words may here be said. The specimens collected during the previous journey as well as those from the two localities first mentioned above (A: 144 etc. and B: 46) HEDIN has called "tuffs", while the other more compact rocks have received various names not indicating a comparatively recent volcanic origin. The short descriptions of their occurrence given in HEDIN's diary never give any indication, suggesting volcanoes with their form still preserved (but compare A: 107). HEDIN also repeatedly tells us that he never has seen a true volcano here. As however the petrographical examination of the rocks collected shows volcanic rocks of modern types to occur here, the conclusion must be drawn, that the volcanic activity that produced them is not

of quite recent date and that therefore the erosion has had time enough to destroy the form of the originally existing volcanoes. On the other hand BONVALOT¹ as well as LITTLEDALE² positively say that they have seen volcanoes in these same regions although only at a distance. LITTLEDALE tells us for instance: "On climbing a ridge to the south — I saw in front seven extinct volcanoes; three were small, but four of them were grand fellows" (p. 461) and: "Between 36° 50' and 33° 50' N. L. (and 87°—88° East Gr.) our path lay through a very volcanic region, numerous undoubted volcanoes being visible" (p. 464). — It appears desirable, that future expeditions to these parts of Eastern Tibet, in spite of the enormous difficulties arising from the inhospitable nature of the country, should nevertheless find time to investigate some of these supposed volcanoes.

Catalogue of the specimens.

A. From June 1900 to March 1901.

1900.

- ³/₇ 1. Greenstone-schist.
- 2. Slate, black, hard, calcareous.
- 3. Granite, quartz-monzonitic.
- ⁴/₇ 3. Dolomite, crystalline, brownish-grey.
- 5. Slate, black, phyllitic.
- 6. » » » , calcareous.
- 7. » » » »
- 8. Granite, red, medium-grained.
- 9. Calcedony-dolomite-rock, brecciated, reddish.
- 10. Greenstone-schist (dynamometamorphic diabase?).
- 11. » » »
- 12. Black siliceous schist.
- 13. Greenstone-schist = 10.
- 14. »
- ⁵/₇ 15. Quartz-felspar-rock, quite crushed.
- 16. Quartz-diorite, schistose.
- 17. Vein-quartz.
- ⁶/₇ 18. Granite, dynamometamorphic.
- 19. Black, hard, phyllitic slate.
- 20. Dark » » » wrinkled up.
- 21. » » » » »
- ⁸/₇ 22. Calcite (from veins).

1900.

- ⁸/₇ 23. Dolomite, dark, close-grained.
- 24. Limestone, dark, impure, close-grained.
- 25. » » » »
- ⁹/₇ 26. Granite, grey fine-grained.
- 27. Greywacke, coarse-grained and fine-grained.
- 28. Greywacke with fragments of black slate.
- 29. Dynamometamorphic greenstone.
- ¹³/₇ 30. Dark, contact-metamorphosized, spotted slate.
- 31. Granite, lightly red, rather coarse-grained.
- 32. Aplite, as dykes in granite.
- ¹⁷/₇ 33. Hornblende-gneiss (containing dykes and veins of red granite).
- 34. Boulder of somewhat crushed plagioclase-granite.
- ²⁰/₇ 35. Coarse red granite.
- 36. Dark, hard hornfels, connected with the granites 35 and 37.
- 37. Coarse red granite = 35.
- 38. Limestone, dense, dark.
- 39. Granite, greenish, schistose.

¹ GABRIEL BONVALOT, De Paris à Tonkin à travers le Tibet inconnu. Paris 1892, p. 188 and the following.

² ST. GEORGE R. LITTLEDALE, A journey across Tibet, from North to South and West to Ladak. (The Geographical Journal Vol. VII No. 5. May 1896.)

1900.

40. Granite, very schistose.
 41. Greywacke, schistose.
^{22/7} 42. » »
 43. Hornblendic granite.
^{23/7} 44. » »
 45. Greywacke-slate, phyllitic.
^{23/7} 46. Greywacke, schistose.
 47. Granite, coarse-grained, light, with two micas.
^{24/7} 48. Granite, fine-grained, grey, with two micas, schistose.
 49. Gneissic granite, connected with 50.
 50. Granite, coarse, grey.
 51. Quartzporphyry, schistose.
 52 a). Quartzporphyry-tuff, schistose.
 b). Granite = type 49 or 50, schistose.
 53. Quartzporphyry-tuff, very fresh, light-yellow with veins of limonite.
 54. Quartzporphyry-tuff, light, compact.
 55. Porphyry or -tuff, much weathered.
^{25/7} 56. Diabase-porphyrityte, fine-grained, weathered; dyke.
 57. Quartz-porphyrityte-tuff.
 58. »
 59. »
 60. »
 61. »
^{26/7} 62. »
 63. Granite, granophyric, with idiomorphic quartz, unaltered, reddish.
 64. Quartzporphyryite-tuff, greenish grey, weathered.
^{31/7} 65. Granite, coarse-grained, grey.
 66. » medium-grained.
 67. » = 65.
^{1/8} 68. » medium-grained, with two micas.
 69. Hornfels, rich in biotite, = 207 and 219.
 70. Granite = 65.
^{3/8} 71. Phyllite.
 72. Sandstone, brownish-red, somewhat schistose.
 73. Phyllite.
 74. Greenstone-schist, probably altered diabase or tuff belonging to the phyllite series.
 75. Limestone, dark-grey, dense.
 76. » » »
 77. » » »
 78. Greenstone, porphyritic, very fine-grained, epidotized.

1900.

- ^{4/8} 79 a). Greywacke, dark.
 b). Greywacke-slate, dark, phyllitic.
 80. Boulders of weathered greywackes.
^{5/8} 81. Greywacke-slate, dark.
 82. Greywacke, comparatively coarse-grained.
^{6/8} 83. » schistose.
 84. » »
 85. » »
 86. » »
^{8/8} 87. » » whitened.
 88. » » »
 89 a). Phyllite, dark-grey.
 b). Greywacke, schistose, whitened.
 90. » » »
 91 a). » » »
 b). Quartzite, rich in calcite, yellow-white, in alternate bedding with a.
^{9/8} 92. Vein-quartz.
 93. Greywacke.
 94. } Volcanic slag, probably andesitic;
 95. } boulders.
^{12/8} 96. Gypsum.
^{14/8} 97. Greywacke.
 98. Greywacke-slate, dark.
^{17/8} 99. Clay slate, black, phyllitic.
 100. Greywacke, greywacke-slate and vein-quartz.
 101. Greywacke.
 102. Conglomerate.
 103 a). Red sandstone.
 b). Conglomerate.
 104. Gypsum.
^{19/8} 105. »
 106. Greywacke, yellow-brown.
^{20/8} 107. Hornblende-andesite.
 108. Dolomite, finely crystalline, yellow; as boulders.
^{22/8} 109. Salt.
 110. = 108, with concentric shell arrangement on a large scale.
^{25/8} 111. Greywacke.
 112. Greywacke-slate with accumulations of carbonaceous matter.
^{27/8} 113. Sandstone, light.
 114. Red sandstone, conglomeratic.
 115. » , more fine-grained.
^{28/8} 116. Limestone, dense, grey (boulder).
 117. » » »
^{30/8} 118. » » »
^{31/8} 119. Porphyryite, weathered, probably dyke.

1900.

- ³¹/₈ 120. Sandstone = 113.
 121. Andesite, weathered.
 122. Red conglomerate.
¹/₉ 123. Red sandstone.
²/₉ 124. »
³/₉ 125. Limestone, dense, grey.
 126. Andesite-tuff, hard and compact.
 127. Red sandstone.
 128. Flint, as boulders.
 129. Windworn black stone.
⁸/₉ 130. Hornblende-andesite, biotite-bearing;
 fresh and weathered specimens.
⁹/₉ 131. Sandstone, greenish.
 132. Red sandstone.
¹⁰/₉ 133. Limestone, dense, dark, bituminous.
 134. » » » »
 135. Hornblende-andesite.
 136. Limestone, dense, dark, bituminous.
 137. » » » »
¹¹/₉ 138. Sandstone, greenish-grey, = 131.
 139. Limestone, dense, dark, bituminous.
¹³/₉ 140. Conglomerate. (As numerous boulders.)
 141. Red sandstone.
 142. Limestone with fossils (?)
 143. Greywacke.
¹⁵/₉ 144. Bronzite-andesite, slaggy, as numerous
 boulders.
¹⁷/₉ 145. » compact, as bed
 rock.
 146. » = 144, boulders.
 147. » compact with some
 vesicles; as bed
 rock.
 148. » = 144, boulders.
 149. » = » »
 150. » = » »
 151. » = » »
 152. » compact, as bed
 rock.
 153. Greywacke-sandstone.
 154. Greywacke, comparatively coarse-
 grained.
 155. Greywacke-sandstone = 153.
¹⁹/₉ 156. Limestone, dense, grey.
²⁰/₉ 157. Greywacke-sandstone, schistose.
 158. Quartzite-sandstone, greyish-white.
 159. » »
²⁴/₉ 160. Red sandstone.
 161. Conglomerate.
²⁵/₉ 162. Red sandstone, very fine-grained.

1900.

- ²⁵/₉ 163. Gypsum.
 164. »
 165. Greywacke-sandstone.
 166. Greywacke-slate, dark.
 167. Limestone, dense, yellow-white.
 168. Greywacke-slate, dark.
²⁷/₉ 169. Porphyry, somewhat schistose and
 much weathered.
 170. Greywacke-slate, close-grained, dark,
 with glistening accumulations of
 carbonaceous matter.
 171. Greywacke-sandstone.
 172. Greywacke.
 173. Conglomerate (belonging to the grey-
 wacke-series?).
 174. Greywacke.
 175. »
 176. Vein-quartz.
²⁸/₉ 177. Limestone, crystalline, black.
 178. Quartz-porphyry, rich in pheno-
 crystals, red.
 179. Quartz-porphyry, rich in pheno-
 crystals, grey.
 180. Quartz-porphyry, rich in pheno-
 crystals, red, weathered.
 181. Granite-porphyry, granophyric, grey,
 connected with the preceding.
 182. Red sandstone.
 183. Greywacke.
 184. »
 185. Greywacke-slate, dark, }
 with carbonaceous ac- } In alternate
 cumulations. } bedding.
 186. Clay-slate, soft, black. }
 187. Greywacke-sandstone.
³⁰/₉ 188. »
 189. Breccia of greywacke and dark lime-
 stone.
 190. Greywacke.
¹/₁₀ 191. Red sandstone.
²/₁₀ 192. Siliceous schist (= B: 41).
 193. Greywacke-slate, phyllitic.
³/₁₀ 194. Greywacke-sandstone, with vein of
 quartz.
 195. Limestone, dense, with dark crystals
 of calcite.
 196. Phyllite.
⁴/₁₀ 197. Limestone, schistose, semicrystalline,
 light.
⁵/₁₀ 198. Conglomerate.
⁶/₁₀ 199. Greywacke.

1900.

200. Conglomerate = 198, connected with:
 201. Calcareous sandstone, fine-grained, light (Greywacke-series?).
 202. Granite, like 215; boulder.
 203. Weathered porphyry-tuff?
 204. Syenite-porphry.
^{6/10} 205. Weathered tuff = 203?
^{7/10} 206. Granite, fine-grained, red, as boulders.
 207. Hornfels (?).
 208. Granite, coarse-grained, with perthitic felspar.
^{8/10} 209. Granite, fine-grained, with two micas, grey.
 210. Mica schist, boulder.
 211. Quartzporphyrite.
 212. Porphyrite, dark, close-grained.
 213. Quartzporphyrite-tuff (boulder).
 214. »
 215. Granite, medium-grained, reddish, not affected by pressure.
 216. Granite, medium-grained, reddish, not affected by pressure.
 217. Granite, = 215 and 216 with aplitic vein.
 218. Greywacke.
 219. »
^{9/10} 220. »
 221. »
 222. »
 223. Clay slate, subordinate layers in 222.
^{10/10} 224. Greywacke.
 225. »
 226. » , schistose.
^{12/10} 227 a). » , almost phylitic.
 b). Quartz-epidote-amphibolite.
^{13/10} 228. Granite with felspar eyes, grey.
 229. Diabase.
 230. Granite-gneiss, red (boulders).
 231. Pegmatite. »
 232. Gneiss, fine-grained, rich in mica.
^{17/10} 233. Granite = 228, intruded in:
 234. Mica schist.
 235. Granite, medium-grained, red.
 236. Granite = 235, vein in red gneiss. (Boulders.)
 237. Granite = 233. (Boulders.)
^{19/10} 238. Green chloritic schist.

1900.

- ^{20/10} 239. Quartzite, bluish white.
^{6/11} 240. Granite, light, somewhat schistose.
 241. » » »
 242. » , gneissic, grey.
 243. » » »
 244. Greywacke.
 245. Quartz-porphryite-tuff.
^{6/11} 246. »
 247. »
 248. »
 249. Greywacke-slate.
 250. Quartz-porphryite-tuff, containing also fragments of dark schists.
 251. Dolomite, crystalline, white.
 252. » » »
 253. Mica schist, coarse.
 254. Quartz-diorite, fine-grained, somewhat schistose.
 255. Granite, gneissic, red.
 256. Dolomite, crystalline, white.
 257. Greywacke, schistose.
 258. Limestone, impure, semicrystalline.
 259. » , schistose.
 260. » »
 261. Granite, dark, rich in hornblende.
 262. Diorite.
 263. Diorite, fine-grained.
 264. » »
 265. Granite, greenish, crushed.
^{11/11} 266. » medium-grained, reddish.
^{12/11} 267. » » , light.
 268. » » , » reddish; rich in microcline, poor in dark minerals.
 269. Diorite, fine-grained = 264.
 270. Quartz-diorite, fine-grained, fresh.
 271. Limestone, impure, dark; with scapolite and diopside.
 272. Calcite-diopside-rock, fine-grained, white.
 273. Limestone, dark = 271.
 274. » black, schistose.
^{13/11} 275. Granite, reddish, medium-grained, with mortar-structure.
 276. Granite, reddish, coarse-grained.
^{14/11} 277. » grey, schistose.
 278. » » »
 279. » » »

1900.

- ¹⁵/₁₁ 280. Granite, grey, schistose.
 281. Granite, red, medium-grained, without strain-phenomena, with perthitic felspar.
 282. Diabase-porphyrity, numerous dykes in:
 283. Granite = 281.
 284. Vein-quartz.
²²/₁₁ 285. Amphibolite, fine-grained.
²²/₁₁ 286. Granite, coarse-grained.
 287. » »
²³/₁₁ 288. » »
 289. Granite, medium-grained, not crushed; with felspar eyes.
 290. Granite, medium-grained, not crushed.
 291. Granite, medium-grained, with two micas, reddish.
 292. Hornfels, dark, rich in brown biotite; perhaps altered greywacke.
²⁹/₁₁ 293. Granite, grey.
 294. Red sandstone.
 295. Granite, schistose, greenish.
 296. » , medium-grained, grey.
 297. Limestone, dark, semicrystalline.
 298. Granite, lenticular, very schistose.
 299. Sericite-schist, perhaps variety of the preceding.
³⁰/₁₁ 300. Dolomite-marble, reddish-white.
 301. » »
 302. Mica schist, calcite- and tourmaline-bearing.
 303. Mica schist, gneissic.
¹/₁₂ 304. » »
 305. » » , light.
³/₁₂ 306. Dolomite-marble, red and white.
 307. » pure white.
 308. Greywacke, dark, fine-grained.
 309. Greywacke, light, coarse-grained, of granitic and dark schist-material.
⁴/₁₂ 310. Greywacke = 308.
 311. Greywacke-slate, dark.
 312. » »
 313. » »
¹³/₁₂ 314. Limestone, dark grey, porous.
¹⁷/₁₂ 315. Sandstone, yellowish, carbonaceous, soft.
 316. Sandstone, yellowish, carbonaceous, soft.
 317. Sandstone, yellowish, carbonaceous, very soft.

1900.

- ¹⁸/₁₂ 318 Sandstone, yellowish, carbonaceous, very soft.
 319. Sandstone, yellowish, carbonaceous, hard, as boulder.
²⁰/₁₂ 320. Granite, crushed and weathered.
 321. » » »
²²/₁₂ 322. Amphibolite.
 323. Greywacke, granite-like, = 309.
 324. Greenstone.
²³/₁₂ 325. Gypsum, crystals.
²⁵/₁₂ 326. Sandstone.
 327. Granite, quartz-monzonitic.
 328. Greywacke, brown, comparatively coarse.
²⁹/₁₂ 329. Greywacke, schistose.
 330. » »
 331. » »
 332. Granite, red, poor in dark minerals.
³⁰/₁₂ 333. Limestone, dense, dark.
 334. » , dense, light with dark streaks.
³¹/₁₂ 335. » , impure, schistose, dark.
 336. » = 333.

1901.

- ¹/₁ 337. Gabbro, schistose.
 338. Limestone, dense, dark grey.
³/₁ 339. Actinolite-schist, close-grained.
 340. Limestone, crystalline, with dark streaks.
 341. Quartz-tourmaline-sericite-schist.
⁴/₁ 342. Greenstone-schist.
 343. » , with calcite veins.
 344. Gabbro with light coloured vein, schistose.
 345. Limestone, crystalline, white.
 346. Granite, crushed and weathered.
 347. Limestone, crystalline, with graphite veins.
 348. Quartz-lense from mica schist.
 349. Sericite-schist.
 350. Serpentine, yellow-green; as boulder.
⁵/₁ 351. Granite, fine-grained, grey, schistose.
 352. Gneiss, fine-grained, grey.
 353. Vein-quartz.
 354. Gneiss, banded.
 355. Granite-gneiss, hornblendic, very schistose, grey.
⁶/₁ 356. Granite-gneiss, hornblendic, very schistose, grey.
¹²/₁ 357. Dolomite-marble.

1901.

- ¹²/₁ 358. Greenstone, porphyritic, dark, decomposed; with dolomite-veins.
- ¹³/₁ 359. Granite, medium-grained, grey.
360. Gneiss, lenticular, grey.
- ¹⁴/₁ 361. Amphibolite.
362. " = schistose gabbro?
363. Muscovite-chlorite-schist.
364. Gneiss, close-grained, grey.
365. Amphibolite.
- ²¹/₁ 366. Gneiss, very rich in hornblende.
- ²²/₁ 367. " "
- ²²/₁ 368. Granulite, yellow-white.
369. Gneiss, garnet-bearing.
370. Mica schist, garnet-bearing.
371. Zoisite-quartz-rock, white.
372. Quartz-amphibolite.
373. Granulite, greyish white.
374. Vein-quartz.
- ²³/₁ 375. Granite-gneiss.
376. Gneiss, dark, garnet-bearing.
377. Dolomite-marble.
- ²⁴/₆ 378. Missing.
- ²⁷/₁ 379. Dolomite-marble.
380. Quartzite.
381. Calcedony-dolomite-rock.
- ²⁸/₁ 382. Granite in contact with kersantite.
383. " " "
384. Granite-gneiss.
385. "
386. "
387. "
388. "
389. "
- ²⁹/₁ 390. "
391. "
392. "
393. "
394. "
395. Granite, coarse-grained, reddish.
396. Limestone, crystalline, white.
- ³⁰/₁ 397. Granite-gneiss.
- ³¹/₁ 398 }
to } Wind-worn stones from the desert.
- ¹/₂ 408. }
- ⁸/₂ 409. Salt, impure, porous.
410. Wind-worn stone.
411. " "
- ⁹/₂ 412. Andesite or porphyrite, basic, weathered.
413. = 412.

1901.

- ⁹/₂ 414. Aplitic granite, red, fine-grained, weathered.
415. = 412.
416. = 412.
417. Trachyte or syenite-porphyry, dense, red, with trachytic structure, somewhat weathered.
418. = 412.
419. = 412.
- ¹⁰/₂ 420. Siliceous schist, dense, dark.
421. Quartz-diabase, medium-grained, showing beginning uralitization.
422. Granophyre.
423. Granite, red.
424. Hornblende-schist.
425. Granite or aplite, white, schistose.
426. " " "
427. Hornblende-schist.
428. "
- ¹¹/₂ 429. Gabbro, schistose.
430. Limestone, semicrystalline, grey.
431. Greywacke, quartzitic.
432. Granite, grey.
433. " , red, even-grained, with dykes of:
434. Diabase, close-grained.
435. Limestone, crystalline.
436. " "
437. " "
438. Amphibolite as dykes, apparently transformed diabase.
439. Granite-gneiss, grey, schistose.
- ¹²/₂ 440. Amphibolite, very schistose.
441. Limestone, finely crystalline, red.
442. Amphibolite, very schistose.
443. Limestone, crystalline, white.
- ¹³/₂ 444. Limestone, crystalline, white, with dark streaks.
445. Greenstone-schist.
- ¹⁴/₂ 446. Chlorite-schist, calcareous, soft.
447. Granite-gneiss, dark.
448. " light.
449. Limestone, crystalline, white.
450. Hornfels or close-grained gneiss, rich in brown biotite.
451. Granite, grey.
452. " red.
- ¹⁶/₂ 453. " dark, schistose.
454. " red.
455. Diabase, transformed into amphibolite.

1901.

- ¹⁶/₃ 456. Granite, crushed, rich in epidote.
 457. = 455.
 458. Quartzite.
 459. Gypsum, crystallized.
¹⁷/₂ 460. Limestone, finely crystalline, schistose, light.
¹⁸/₂ 461. Diabase.
 462. Dolomite, crystalline, grey.
 463. » » »

1901.

- ¹⁸/₂ 464. Limestone = 460.
 465. Greywacke.
 466. »
²³/₂ 467. »
 468. Limestone, schistose, dense, yellow; boulder.
²⁴/₂ 469. Greywacke.
¹/₃ 470. »
 471. »

B. From May to December 1901.

1901.

- ¹⁸/₅ 1. Gneiss with pegmatitic veins.
 2. Gneiss, fine-grained, rich in biotite.
 3 a). » » »
 b). Dolomite.
 4. Gneiss = 2.
 5. Dolomite, white, coarsely crystalline.
 6. Granite-gneiss, light.
²⁰/₅ 7. Limestone, grey, soft, fine-grained.
 8. Granite, light, very schistose.
 9. » » »
²¹/₅ 10. » » »
 11. » » »
 12. » pegmatitic, less schistose.
 13. » light, schistose.
²²/₅ 14. Gneiss, fine-grained, hornblende-bearing.
 15. » » »
 16. » » »
²³/₅ 17. Pegmatitic nodule.
 18. Granite-gneiss, grey, rich in plagioclase.
 19. Pyroxene-gneiss with scapolite and calcite.
 20. Granite, red, coarsely crystalline.
 21. Gneiss = 18.
 22. Quartz, apparently from veins.
 23. Granite, red, syenitic.
²⁵/₅ 24. Limestone with fragments of quartzitic rocks.
 25. Gneiss = 2, rich in plagioclase.
 26. Gneiss = 2.
²⁷/₅ 27. Dolomite, dark, fine-grained.
 28. Gneiss, grey, = 26 etc.
²⁸/₅ 29. Red sandstone.
³⁰/₅ 30. Mica schist, rich in tourmaline (conf. A: 341).
 31. Quartzite.

1901.

- ¹/₆ 32. Sericite-schist, light green.
 33. Gneiss, with strongly developed schistosity.
⁶/₆ 34. Gypsum.
 35. »
 36. Sandstone, soft, yellow.
⁸/₆ 37. } Missing; "red clay" according to the
¹⁰/₆ 38. } diary.
¹²/₆ 39. Sandstone.
¹⁴/₆ 40. Greywacke-slate, green, schistose.
¹⁷/₆ 41. Siliceous schist (= A: 192).
¹⁸/₆ 42. Limestone, light-grey, fine-grained.
 43. Greywacke.
²³/₆ 44. »
 45. »
²⁴/₆ 46. Bronzite-andesite, vesicular.
 47. Crystals of quartz, as boulders.
²⁵/₆ 48. Greywacke.
 49. »
 50. Vein-quartz.
 51. Groupe of aragonite needles, radially arranged.
²⁶/₆ 52. Greywacke-sandstone.
²⁸/₆ 53. »
 54. »
 55. Hard siliceous concretions.
²⁹/₆ 56. Diorite-porphyrity, much weathered.
 57. Greywacke.
 58. Black slate.
 59. Quartz-crystal (boulder).
³⁰/₆ 60. Gypsum.
 61. Greywacke.
¹/₇ 62. »
 63. Gypsum, crystallized ("in enormous boulders").
 64. Greywacke.
 65. Black slate (= 58).

1901.

- ¹/₇ 66. Red conglomerate.
 67. » »
 68. Gypsum.
²/₇ 69. Red sandstone.
 70. » »
⁴/₇ 71. Phyllite.
 72. Quartzite-sandstone, grey (= A: 158).
 73. Greenstone-schist, alternating with:
 74. } Siliceous schist, dense, black, with
 75. } calcite veins.
⁸/₇ 76. Oolitic calcareous sandstone.
⁹/₇ 77. Limestone, dense, grey.
¹⁰/₇ 78. Limestone (with fossils?).
 79. Violet-red sandstone.
 80. Limestone, dense, dark, with white veins.
¹¹/₇ 81. Fragments of hard siliceous chert with etchings. (Boulders.)
 82. Conglomerate with gypsum on cavities.
¹²/₇ 83. Limestone, dark, bituminous.
 84. » » »
¹⁸/₇ 85. » » »
¹⁹/₇ 86. Porous calcareous rock.
 87. Sandstone, light-grey.
 88. Magnetic iron ore, porous, as boulders.
²¹/₇ 89. Limestone, dark, bituminous (with fossils?).
 90. Limestone, dark, bituminous.
 91. Calcareous slate.
 92. Limestone, dark.
²²/₇ 93. Calcareous slate = 91.
 94. Conglomerate with fragments of slate (= 101).
²⁴/₈ * 95. Quartz-porphyrity-tuff, rather weathered, but without signs of strain and therefore perhaps "young-volcanic". (Boulders.)
 96. Limestone, dark-grey, with fossils.
 97. Greywacke-slate.
 98. Limestone with fossils.
 99. Sandstone.
 100. Red conglomerate.
 101. Green sandstone with pebbles of limestone and black slate.
²⁵/₈ 102. Sandstone.
 103. Limestone, dark, with fossils.

1901.

- ²⁸/₈ 104. Greywacke-sandstone, hard, greyish-brown.
²⁹/₈ 105. }
 106. } Limestone, dark, dense, bituminous.
 107. }
 108. }
¹/₉ 109. Calcareous sandstone, brick-red, loose, comparatively coarse-grained.
 110. Limestone, grey, dense.
 111. » yellow-white.
³/₉ 112. Greywacke.
⁴/₉ 113. Greenish-grey sandstone, marly, soft.
 114. » » » »
⁵/₉ 115. Sandstone, brownish-grey.
⁷/₉ 116. Limestone, grey, dense, with chert.
⁸/₉ 117. » » » »
 118. Red sandstone.
 119. Conglomerate.
¹⁰/₉ 120. } Limestone, light, dense, fossiliferous,
¹¹/₉ 121. } with chert.
 122. Sandstone.
 123. Bronzite-andesite, holocrystalline.
 124. }
¹⁴/₉ 125. }
 126. } Limestone, light-grey, dense, apparently fossiliferous.
¹⁵/₉ 127. }
¹⁸/₉ 128. }
 129 a. }
²⁵/₉ 129 b. Limestone-conglomerate.
 130. } Limestone, light-grey, dense, with
 131. } fossils.
 132. Red sandstone.
 133. »
²⁸/₉ 134. »
²⁹/₉ 135. Limestone, light, dense.
²/₁₀ 136. » » »
³/₁₀ 137. » » »
 138. » » »
⁴/₁₀ 139. Conglomerate, red, = 119.
 140. Conglomerate (= 94).
 141. Diabase.
 142. » pale, weathered.
 143. » » »
 144. White granophyric dyke rock (quartz-bostonite). (Boulder.)
 145. Light violet-grey dyke rock (quartz-bostonite). (Boulder.)
 146. Black phyllite. (Boulder.)

* The dash for Lhasa. Scientific Results III p. 499—525.

1901.

- ⁴/₁₀ 147. Like 144 and 145. (Boulder.)
 148. Quartzite-sandstone = 168. (Boulder.)
 149. } Like 144, 145 and 147, weathered.
 150. } (Boulder.)
 151. Diabase. (Boulder.)
 152. Weathered calcareous rock.
 153. Hornblende-andesite, holocrystalline, somewhat weathered. (Boulder.)
⁵/₁₀ 154. Porphyrite, much weathered.
 155. " "
⁷/₁₀ 156. Quartz, white, granular, probably from veins.
⁸/₁₀ 157. Black phyllite.
¹⁰/₁₀ 158. Limestone, grey, dense, with fossils?
 159. " "
¹¹/₁₀ 160. }
 161. } Limestone, yellowish white, finely crystalline.
¹³/₁₀ 162. }
¹⁵/₁₀ 163. } Limestone, yellowish white, dense,
¹⁶/₁₀ 164. } brecciated.
¹⁸/₁₀ 165. Greywacke, fine-grained.
 166. Calc-sinter, apparently recent.
²⁰/₁₀ 167. Conglomerate, calcareous, probably recent.
 168. Quartzite-sandstone = 148.
²¹/₁₀ 169. Limestone, white, finely crystalline.
 170. Greywacke, dark, fine-grained.
 171. " "
²²/₁₀ 172. Dolomite, dense, yellowish white.
²³/₁₀ 173. Quartzite-sandstone.
 174. "
²⁴/₁₀ 175. Diabase, probably dyke in the following.
 176. Limestone, white, dense.
²⁵/₁₀ 177. " pink, finely crystalline.
²⁶/₁₀ 178. " yellowish-grey, dense.
²⁷/₁₀ 179. Rhyolite-tuff, hard, compact.
 180. Trachyte.
 181. "
 182. Rhyolite-tuff.
 183. Trachyte with veins of agate.
 184. "
 185. = 183.
 186. Rhyolite-tuff, hard, compact.
 187. " "
²⁸/₁₀ 188. Boulder of chalcopyrite, malachite and azurite.
 189. } Greenish grey phyllite or greywacke-
 190. } schist.
 191. Granophyric granite, boulder.

1901.

- ²⁸/₁₀ 192. Granite, aplitic dyke in the schist 189—190.
 193. Porphyrite, grey, unaltered, fine-grained.
 194. Granite, type: "Quartz-monzonite", medium-grained, very fresh.
 195. Porphyrite, dark, basic; dyke in the foregoing.
 196. Porphyrite, dark, basic.
³/₁₁ 197. Calc-sinter, recent. (Boulder.)
⁶/₁₁ 198. Porphyrite or -tuff, decomposed.
⁷/₁₁ 199. " "
 200. Porphyrite or -tuff, decomposed, brecciated.
 201. Porphyrite or -tuff, decomposed.
 202. Calc-sinter, with enclosed sand grains.
⁹/₁₁ 203. Conglomerate, red, with calcareous matrix.
¹¹/₁₁ 204. Limestone, grey, dense.
¹³/₁₁ 205. Limestone, grey, dense.
¹⁷/₁₁ 206. }
 207. } Limestone, reddish-white, finely crystalline.
 208. Porphyrite, decomposed.
 209. Limestone, white, finely crystalline.
 210. Black phyllite (boulder).
¹⁸/₁₁ 211. Limestone, dense, yellowish grey.
¹⁹/₁₁ 212. Limestone, finely crystalline, reddish white.
 213. Limestone, finely crystalline, white.
 214. Limestone, finely crystalline, reddish white.
 215. Limestone, finely crystalline, reddish white.
 216. Phyllite, hard.
²⁰/₁₁ 217. Limestone, dark, bituminous, finely crystalline.
²⁵/₁₁ 218. Missing.
 219. Limestone, dense, dark.
²⁶/₁₁ 220. " "
²⁷/₁₁ 221. " "
 222. " "
²⁸/₁₁ 223. " " , brecciated.
 224. Limestone, finely crystalline, white.
³⁰/₁₁ 225. " " , grey.
 226. " "
 227. Very schistose porphyry (?), decomposed.
 228. Calcite, probably from veins.
 229. Limestone, finely crystalline, grey.
¹/₁₂ 230. " dense, dark.
 231. Calcite = 228.

1901.

- ¹/₁₂ 232. Calcite = 228.
 233. Porphyrite-tuff.
 234. Green phyllitic schist in alternate bedding with:
 235. Limestone, finely crystalline, brownish.
 236. Limestone, crystalline, white.
²/₁₂ 237. Calcareous slate.
 238. Limestone, dark.
 239. Biotite-plagioclase-granite, fresh.
 240. Phyllite, black.
 241. Greywacke, dark, phyllitic.
 242. Phyllite.
³/₁₂ 243. »
 244. »
⁴/₁₂ 245. » (like 241).
 246. »
¹⁰/₁₂ 247. Greenstone-schist, apparently connected with the phyllites.
¹⁰/₁₂ 248. Sericite-schist, apparently connected with the phyllites.
 249. Talc-schist with siderite.
 250. » » quartz veins.
¹¹/₁₂ 251. Calc-sinter-conglomerate, recent.
 252. Greenstone-schist = 247.
 253. Phyllite.
 254. Very schistose quartz-porphyry.
¹²/₁₂ 255. »
 256. »
 257. »
 258. »
 259. » decomposed.

1901.

- ¹²/₁₂ 260. Talc-schist, with siderite.
 261. Weathered yellow rock.
¹³/₁₂ 262. Phyllite.
 263. Vein-quartz.
¹⁴/₁₂ 264. Calc-sinter.
 265. Greywacke-slate, hard, dark grey.
 266. Limestone, crystalline, white.
 267. Granite, medium-grained, grey, fresh.
 268. Greywacke-slate = 265.
¹⁴/₁₂ 269. Phyllite, dark-green.
 270. » »
 271. Weathered schist.
¹⁵/₁₂ 272. Mica-schist, dark, fine-grained, with garnet.
 273. Dolomite, finely crystalline, schistose.
 274. Vein-quartz, red (boulder).
¹⁶/₁₂ 275. Mica-schist, dark, fine-grained.
 276. Vein-quartz.
 277. Quartz-monzonite = 194 (boulder).
 278. Dolomite, crystalline, white.
 279. Limestone, coarsely crystalline, white.
 280. Dolomite, crystalline, white.
 281. Phyllite.
 282. Hornblende-schist (boulder).
¹⁷/₁₂ 283. Granite (white soda-granite).
 284. » »
 285. » » gneissic.
 286. Hornblende-biotite-rock, apparently basic inclusion in the preceding.
 287. Mica schist, hornblende-bearing.

Profile from Leh to Kargalik 1902.

1902.

- ¹¹/₄ a. Granite, schistose, white; »soda-granite».
 b. » » , reddish grey, »
 c. » » , grey; »
¹³/₄ d. » » » »
¹⁴/₄ e. » » , white; »
¹⁵/₄ f. Dolomite, grey, almost dense.
 g. Dolomite, white, crystalline.
¹⁶/₄ h. Hornblende-granite.
¹⁷/₄ i. Granite, grey, = B: 267.
¹⁸/₄ j. Quartz-diorite, fine-grained.
 k. Greywacke-slate, hard, black.
¹⁰/₄ l. Granite, light, weathered.
 m. Actinolite-zoisite-quartz-rock, close-grained.
 n. Limestone, dense, dark.

1902.

- ¹⁰/₄ o. Limestone, dolomitic, finely crystalline, light-grey.
²¹/₄ p. Calcite, coarsely crystalline, probably from veins.
 q. Dolomite, finely crystalline, white.
 r. Limestone, dense, yellowish.
 s. » » , grey.
²³/₄ t. » » , bituminous, dark.
²⁴/₄ u. Black phyllite.
 v. Limestone, dense, brown.
 w. Limestone, finely crystalline, light grey.
²⁴/₄ x. Limestone, finely crystalline, light grey, crushed.
 y. Limestone, impure, schistose, grey.
²⁶/₄ z. Greywacke (?) schistose, weathered.

1902.

- ²⁶/₄ ä. Phyllite, hard, black.
²⁸/₄ ä. Sericite-schist, probably strongly schistose granite.
²⁹/₄ ö. Gneiss, light grey, probably strongly schistose granite.
³/₅ a'. Quartz, granular, white with dark strings.
 b'. Granite, medium-grained, light grey.
 c'. » , fine-grained, » »
 d'. » , medium-grained, schistose, darker.
²/₅ e'. » , coarse grained, non-schistose, light.

1902.


- ²/₅ f'. Mica-schist or strongly schistose granite.
³/₅ g'. » » »
 h'. Quartz, granular (= a') and black schist.
⁴/₅ i'. Crushed pegmatite or coarsely crystalline granite.
 j'. Greywacke, fine-grained, dark, typical.
⁵/₅ k'. Mica-schist or strongly schistose granite, garnet-bearing.
 l'. Missing.
 m'. »

Geological Map
of the route of Dr. Sven Hedin in Tibet 1899-1902

Helge Backström and Harald Johansson

Scale 1:2000000

0 10 20 30 40 50 60 70 80 90 100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360 370 380 390 400 410 420 430 440 450 460 470 480 490 500 510 520 530 540 550 560 570 580 590 600 610 620 630 640 650 660 670 680 690 700 710 720 730 740 750 760 770 780 790 800 810 820 830 840 850 860 870 880 890 900 910 920 930 940 950 960 970 980 990 1000 1010 1020 1030 1040 1050 1060 1070 1080 1090 1100 1110 1120 1130 1140 1150 1160 1170 1180 1190 1200 1210 1220 1230 1240 1250 1260 1270 1280 1290 1300 1310 1320 1330 1340 1350 1360 1370 1380 1390 1400 1410 1420 1430 1440 1450 1460 1470 1480 1490 1500 1510 1520 1530 1540 1550 1560 1570 1580 1590 1600 1610 1620 1630 1640 1650 1660 1670 1680 1690 1700 1710 1720 1730 1740 1750 1760 1770 1780 1790 1800 1810 1820 1830 1840 1850 1860 1870 1880 1890 1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000 2010 2020 2030 2040 2050 2060 2070 2080 2090 2100 2110 2120 2130 2140 2150 2160 2170 2180 2190 2200 2210 2220 2230 2240 2250 2260 2270 2280 2290 2300 2310 2320 2330 2340 2350 2360 2370 2380 2390 2400 2410 2420 2430 2440 2450 2460 2470 2480 2490 2500 2510 2520 2530 2540 2550 2560 2570 2580 2590 2600 2610 2620 2630 2640 2650 2660 2670 2680 2690 2700 2710 2720 2730 2740 2750 2760 2770 2780 2790 2800 2810 2820 2830 2840 2850 2860 2870 2880 2890 2900 2910 2920 2930 2940 2950 2960 2970 2980 2990 3000 3010 3020 3030 3040 3050 3060 3070 3080 3090 3100 3110 3120 3130 3140 3150 3160 3170 3180 3190 3200 3210 3220 3230 3240 3250 3260 3270 3280 3290 3300 3310 3320 3330 3340 3350 3360 3370 3380 3390 3400 3410 3420 3430 3440 3450 3460 3470 3480 3490 3500 3510 3520 3530 3540 3550 3560 3570 3580 3590 3600 3610 3620 3630 3640 3650 3660 3670 3680 3690 3700 3710 3720 3730 3740 3750 3760 3770 3780 3790 3800 3810 3820 3830 3840 3850 3860 3870 3880 3890 3900 3910 3920 3930 3940 3950 3960 3970 3980 3990 4000 4010 4020 4030 4040 4050 4060 4070 4080 4090 4100 4110 4120 4130 4140 4150 4160 4170 4180 4190 4200 4210 4220 4230 4240 4250 4260 4270 4280 4290 4300 4310 4320 4330 4340 4350 4360 4370 4380 4390 4400 4410 4420 4430 4440 4450 4460 4470 4480 4490 4500 4510 4520 4530 4540 4550 4560 4570 4580 4590 4600 4610 4620 4630 4640 4650 4660 4670 4680 4690 4700 4710 4720 4730 4740 4750 4760 4770 4780 4790 4800 4810 4820 4830 4840 4850 4860 4870 4880 4890 4900 4910 4920 4930 4940 4950 4960 4970 4980 4990 5000 5010 5020 5030 5040 5050 5060 5070 5080 5090 5100 5110 5120 5130 5140 5150 5160 5170 5180 5190 5200 5210 5220 5230 5240 5250 5260 5270 5280 5290 5300 5310 5320 5330 5340 5350 5360 5370 5380 5390 5400 5410 5420 5430 5440 5450 5460 5470 5480 5490 5500 5510 5520 5530 5540 5550 5560 5570 5580 5590 5600 5610 5620 5630 5640 5650 5660 5670 5680 5690 5700 5710 5720 5730 5740 5750 5760 5770 5780 5790 5800 5810 5820 5830 5840 5850 5860 5870 5880 5890 5900 5910 5920 5930 5940 5950 5960 5970 5980 5990 6000 6010 6020 6030 6040 6050 6060 6070 6080 6090 6100 6110 6120 6130 6140 6150 6160 6170 6180 6190 6200 6210 6220 6230 6240 6250 6260 6270 6280 6290 6300 6310 6320 6330 6340 6350 6360 6370 6380 6390 6400 6410 6420 6430 6440 6450 6460 6470 6480 6490 6500 6510 6520 6530 6540 6550 6560 6570 6580 6590 6600 6610 6620 6630 6640 6650 6660 6670 6680 6690 6700 6710 6720 6730 6740 6750 6760 6770 6780 6790 6800 6810 6820 6830 6840 6850 6860 6870 6880 6890 6900 6910 6920 6930 6940 6950 6960 6970 6980 6990 7000 7010 7020 7030 7040 7050 7060 7070 7080 7090 7100 7110 7120 7130 7140 7150 7160 7170 7180 7190 7200 7210 7220 7230 7240 7250 7260 7270 7280 7290 7300 7310 7320 7330 7340 7350 7360 7370 7380 7390 7400 7410 7420 7430 7440 7450 7460 7470 7480 7490 7500 7510 7520 7530 7540 7550 7560 7570 7580 7590 7600 7610 7620 7630 7640 7650 7660 7670 7680 7690 7700 7710 7720 7730 7740 7750 7760 7770 7780 7790 7800 7810 7820 7830 7840 7850 7860 7870 7880 7890 7900 7910 7920 7930 7940 7950 7960 7970 7980 7990 8000 8010 8020 8030 8040 8050 8060 8070 8080 8090 8100 8110 8120 8130 8140 8150 8160 8170 8180 8190 8200 8210 8220 8230 8240 8250 8260 8270 8280 8290 8300 8310 8320 8330 8340 8350 8360 8370 8380 8390 8400 8410 8420 8430 8440 8450 8460 8470 8480 8490 8500 8510 8520 8530 8540 8550 8560 8570 8580 8590 8600 8610 8620 8630 8640 8650 8660 8670 8680 8690 8700 8710 8720 8730 8740 8750 8760 8770 8780 8790 8800 8810 8820 8830 8840 8850 8860 8870 8880 8890 8900 8910 8920 8930 8940 8950 8960 8970 8980 8990 9000 9010 9020 9030 9040 9050 9060 9070 9080 9090 9100 9110 9120 9130 9140 9150 9160 9170 9180 9190 9200 9210 9220 9230 9240 9250 9260 9270 9280 9290 9300 9310 9320 9330 9340 9350 9360 9370 9380 9390 9400 9410 9420 9430 9440 9450 9460 9470 9480 9490 9500 9510 9520 9530 9540 9550 9560 9570 9580 9590 9600 9610 9620 9630 9640 9650 9660 9670 9680 9690 9700 9710 9720 9730 9740 9750 9760 9770 9780 9790 9800 9810 9820 9830 9840 9850 9860 9870 9880 9890 9900 9910 9920 9930 9940 9950 9960 9970 9980 9990 10000 10010 10020 10030 10040 10050 10060 10070 10080 10090 10100 10110 10120 10130 10140 10150 10160 10170 10180 10190 10200 10210 10220 10230 10240 10250 10260 10270 10280 10290 10300 10310 10320 10330 10340 10350 10360 10370 10380 10390 10400 10410 10420 10430 10440 10450 10460 10470 10480 10490 10500 10510 10520 10530 10540 10550 10560 10570 10580 10590 10600 10610 10620 10630 10640 10650 10660 10670 10680 10690 10700 10710 10720 10730 10740 10750 10760 10770 10780 10790 10800 10810 10820 10830 10840 10850 10860 10870 10880 10890 10900 10910 10920 10930 10940 10950 10960 10970 10980 10990 11000 11010 11020 11030 11040 11050 11060 11070 11080 11090 11100 11110 11120 11130 11140 11150 11160 11170 11180 11190 11200 11210 11220 11230 11240 11250 11260 11270 11280 11290 11300 11310 11320 11330 11340 11350 11360 11370 11380 11390 11400 11410 11420 11430 11440 11450 11460 11470 11480 11490 11500 11510 11520 11530 11540 11550 11560 11570 11580 11590 11600 11610 11620 11630 11640 11650 11660 11670 11680 11690 11700 11710 11720 11730 11740 11750 11760 11770 11780 11790 11800 11810 11820 11830 11840 11850 11860 11870 11880 11890 11900 11910 11920 11930 11940 11950 11960 11970 11980 11990 12000 12010 12020 12030 12040 12050 12060 12070 12080 12090 12100 12110 12120 12130 12140 12150 12160 12170 12180 12190 12200 12210 12220 12230 12240 12250 12260 12270 12280 12290 12300 12310 12320 12330 12340 12350 12360 12370 12380 12390 12400 12410 12420 12430 12440 12450 12460 12470 12480 12490 12500 12510 12520 12530 12540 12550 12560 12570 12580 12590 12600 12610 12620 12630 12640 12650 12660 12670 12680 12690 12700 12710 12720 12730 12740 12750 12760 12770 12780 12790 12800 12810 12820 12830 12840 12850 12860 12870 12880 12890 12900 12910 12920 12930 12940 12950 12960 12970 12980 12990 13000 13010 13020 13030 13040 13050 13060 13070 13080 13090 13100 13110 13120 13130 13140 13150 13160 13170 13180 13190 13200 13210 13220 13230 13240 13250 13260 13270 13280 13290 13300 13310 13320 13330 13340 13350 13360 13370 13380 13390 13400 13410 13420 13430 13440 13450 13460 13470 13480 13490 13500 13510 13520 13530 13540 13550 13560 13570 13580 13590 13600 13610 13620 13630 13640 13650 13660 13670 13680 13690 13700 13710 13720 13730 13740 13750 13760 13770 13780 13790 13800 13810 13820 13830 13840 13850 13860 13870 13880 13890 13900 13910 13920 13930 13940 13950 13960 13970 13980 13990 14000 14010 14020 14030 14040 14050 14060 14070 14080 14090 14100 14110 14120 14130 14140 14150 14160 14170 14180 14190 14200 14210 14220 14230 14240 14250 14260 14270 14280 14290 14300 14310 14320 14330 14340 14350 14360 14370 14380 14390 14400 14410 14420 14430 14440 14450 14460 14470 14480 14490 14500 14510 14520 14530 14540 14550 14560 14570 14580 14590 14600 14610 14620 14630 14640 14650 14660 14670 14680 14690 14700 14710 14720 14730 14740 14750 14760 14770 14780 14790 14800 14810 14820 14830 14840 14850 14860 14870 14880 14890 14900 14910 14920 14930 14940 14950 14960 14970 14980 14990 15000 15010 15020 15030 15040 15050 15060 15070 15080 15090 15100 15110 15120 15130 15140 15150 15160 15170 15180 15190 15200 15210 15220 15230 15240 15250 15260 15270 15280 15290 15300 15310 15320 15330 15340 15350 15360 15370 15380 15390 15400 15410 15420 15430 15440 15450 15460 15470 15480 15490 15500 15510 15520 15530 15540 15550 15560 15570 15580 15590 15600 15610 15620 15630 15640 15650 15660 15670 15680 15690 15700 15710 15720 15730 15740 15750 15760 15770 15780 15790 15800 15810 15820 15830 15840 15850 15860 15870 15880 15890 15900 15910 15920 15930 15940 15950 15960 15970 15980 15990 16000 16010 16020 16030 16040 16050 16060 16070 16080 16090 16100 16110 16120 16130 16140 16150 16160 16170 16180 16190 16200 16210 16220 16230 16240 16250 16260 16270 16280 16290 16300 16310 16320 16330 16340 16350 16360 16370 16380 16390 16400 16410 16420 16430 16440 16450 16460 16470 16480 16490 16500 16510 16520 16530 16540 16550 16560 16570 16580 16590 16600 16610 16620 16630 16640 16650 16660 16670 16680 16690 16700 16710 16720 16730 16740 16750 16760 16770 16780 16790 16800 16810 16820 16830 16840 16850 16860 16870 16880 16890 16900 16910 16920 16930 16940 16950 16960 16970 16980 16990 17000 17010 17020 17030 17040 17050 17060 17070 17080 17090 17100 17110 17120 17130 17140 17150 17160 17170 17180 17190 17200 17210 17220 17230 17240 17250 17260 17270 17280 17290 17300 17310 17320 17330 17340 17350 17360 17370 17380 17390 17400 17410 17420 17430 17440 17450 17460 17470 17480 17490 17500 17510 17520 17530 17540 17550 17560 17570 17580 17590 17600 17610 17620 17630 17640 17650 17660 17670 17680 17690 17700 17710 17720 17730 17740 17750 17760 17770 17780 17790 17800 17810 17820 17830 17840 17850 17860 17870 17880 17890 17900 17910 17920 17930 17940 17950 17960 17970 17980 17990 18000 18010 18020 18030 18040 18050 18060 18070 18080 18090 18100 18110 18120 18130 18140 18150 18160 18170 18180 18190 18200 18210 18220 18230 18240 18250 18260 18270 18280 18290 18300 18310 18320 18330 18340 18350 18360 18370 18380 18390 18400 18410 18420 18430 18440 18450 18460 18470 18480 18490 18500 18510 18520 18530 18540 18550 18560 18570 18580 18590 18600 18610 18620 18630 18640 18650 18660 18670 18680 18690 18700 18710 18720 18730 18740 18750 18760 18770 18780 18790 18800 18810 18820 18830 18840 18850 18860 18870 18880 18890 18900 18910 18920 18930 18940 18950 18960 18970 18980 18990 19000 19010 19020 19030 19040 19050 19060 19070 19080 19090 19100 19110 19120 19130 19140 19150 19160 19170 19180 19190 19200 19210 19220 19230 19240 19250 19260 19270 19280 19290 19300 19310 19320 19330 19340 19350 19360 19370 19380 19390 19400 19410 19420 19430 19440 19450 19460 19470 19480 19490 19500 19510 19520 19530 19540 19550 19560 19570 19580 19590 19600 19610 19620 19630 19640 19650 19660 19670 19680 19690 19700 19710 19720 19730 19740 19750 19760 19770 19780 19790 19800 19810 19820 19830 19840 19850 19860 19870 19880 19890 19900 19910 19920 19930 19940 19950 19960 19970 19980 19990 20000 20010 20020 20030 20040 20050 20060 20070 20080 20090 20100 20110 20120 20130 20140 20150 20160 20170 20180 20190 20200 20210 20220 20230 20240 20250 20260 20270 20280 20290 20300 20310 20320 20330 20340 20350 20360 20370 20380 20390 20400 20410 20420 20430 20440 20450 20460 20470 20480 20490 20500 20510 20520 20530 20540 20550 20560 20570 20580 20590 20600 20610 20620 20630 20640 20650 20660 20670 20680 20690 20700 20710 20720 20730 20740 20750 20760 20770 20780 20790 20800 20810 20820 20830 20840 20850 20860 20870 20880 20890 20900 20910 20920 20930 20940 20950 20960 20970 20980 20990 21000 21010 21020 21030 21040 21050 21060 21070 21080 21090 21100 21110 21120 21130 21140 21150 21160 21170 21180 21190 21200 21210 21220 21230 21240 21250 21260 21270 21280 21290 21300 21310 21320 21330 21340 21350 21360 21370 21380 21390 21400 21410 21420 21430 21440 21450 21460 21470 21480 21490 21500 21510 21520 21530 21540 21550 21560 21570 21580 21590 21600 21610 21620 21630 21640 21650 21660 21670 21680 21690 21700 21710 21720 21730 21740 21750 21760 21770 21780 21790 21800 21810 21820 21830 21840 21850 21860 21870 21880 21890 21900 21910 21920 21930 21940 21950 21960 21970 21980 21990 22000 22010 22020 22030 22040 22050 22060 22070 22080 22090 22100 22110 22120 22130 22140 22150 22160 22170 22180 22190 2



T

PRESIDENT'S
SECRETARIAT
LIBRARY

T